

有吃有保佑 吃落出代誌

情緒類營養品解析

徐莉寧 (DVM, MSc, Ms)

2020/12/20

今日大綱

- 行為問題的本質與困境
- 為什麼選擇營養補充品(Supplements)
- 補充之前，先談營養
- 成分個論
- 國內外常見產品
- 注意事項



獸醫師國家考試合格

中興大學獸醫系 學士

台灣大學獸醫所 神經生理實驗室 碩士

英國林肯大學臨床動物行為學 碩士

長庚醫學大學 神經內科/生物機電 研究助理

湖光動物醫院 獸醫師

粉專- 獸醫師阿默 小動物的日常行為

會思考的狗工作室 行為獸醫師

Fear Free Professional certificated

現駐診於逗號動物醫院





逗號動物醫院
Dog House Animal Hospital




行為問題的本質與困境

心理性疾患的異質性 (heterogeneous)

- 診斷標準(Criteria) 不停修改
- 各國家/學會發展各自的標準/詞彙 (Term)
- 大量使用問卷 (Questionnaire) 與量表 (Scale)
- 飼主的觀察/ 主觀因素
- 同樣的行為表象，不同的動機
- 同樣的診斷，不同的原因
- 個案間差異太大，很難有統一的建議與處置





Depression:

- Neurotransmitters: Norepinephrine, Serotonin, Dopamine...
- Brain regions: Frontal lobe, Hippocampus, Temporal lobe, Thalamus, Striatum, and Amygdala...
- Imbalance of Monoamine > Serotonin deficiency

Anxiety:

- Neurotransmitters: GABA, Serotonin, Norepinephrine...
- Brain regions: Amygdala, Hippocampus, dorsal anterior cingulate Cortex (dACC), and Ventromedial prefrontal Cortex...

Obsessive-compulsive disorder (OCD)

- Neurotransmitters: Glutamate, GABA, Dopamine...
- Brain regions: Striatum (caudate and putamen), anterior Thalamus, anterior cingulate Cortex, and orbitofrontal Cortex

我們到底在討論什麼？

- Animal behavior (動物行為)
 - Ethology (動物行為) : Ecology, Biology, Evolution, Comparative psychology...
 - Anthropomorphism
 - Animal mind...? : Cognition, Spirit, Consciousness, Intelligence...



為什麼選擇營養補充品 (Supplements)

營養品(Supplements)的優勢

- 對身心科藥物的刻板印象
- 相較於藥物與訓練，風險較低
- 使用限制較少
- 老年、其他醫療考量
- 未完全投予也無妨
- 不穩定/不可靠的飼主



營養品(Supplements)的劣勢

- 效果有限
- 個體差異大，報告結果無定論
- 投予對象常為挑食、食慾不佳
- 多為口嚼錠，若適口性不佳，投予困難
- 成本高昂
- 大部分需要使用一段時間

補充之前，先談營養

Nutraceuticals : 營養醫學/食療



- A balanced mood is supported by a balanced diet.

營養均衡，才有穩定的情緒！以下營養素都參與了情緒調節

- Protein: L-tryptophan, Tyrosine...
- Fat: Cholesterol, Omega-3, Omega-6...
- Carbohydrate: Insulin
- Vitamins: B, C...
- Trace Minerals: Magnesium, Calcium...

單單提個胺基酸....

- L-Tryptophan availability⁽³⁾
 - Large neutral amino acids (LNAA): Tyrosine, Leucine, Isoleucine, Valine, Phenylalanine ...
 - Competitive nature of the transporter for LNAA at the blood-brain barrier.
 - A High L-Tryptophan/LNAAs meal promote L-Tryptophan availability in brain.
 - Carbohydrate ingestion increased the secretion of insulin which raised plasma levels of tryptophan and lowered the concentrations of the competing amino acids.
 - Vitamin B₆ act as rate limiting cofactor in the synthesis of neurotransmitters



Comfort food

From Wikipedia, the free encyclopedia

For the Noah Ashenhurst novel, see [Comfort Food \(novel\)](#).

Comfort food is kinds of [food](#) that provides a [nostalgic](#) or sentimental value to someone,^[1] and may be characterized by its high [caloric nature](#), high [carbohydrate](#) level, or simple [preparation](#).^[2] The nostalgia may be specific to an individual, or it may apply to a specific [culture](#).^[3]

Contents [hide]

- 1 Definition
- 2 Psychological studies
- 3 By region
 - 3.1 Afghanistan
 - 3.2 Australia and New Zealand
 - 3.3 Canada
 - 3.4 Egypt
 - 3.5 France
 - 3.6 Hong Kong
 - 3.7 India
 - 3.8 Indonesia
 - 3.9 Italy
 - 3.10 Japan
 - 3.11 Philippines
 - 3.12 Poland
 - 3.13 Puerto Rico
 - 3.14 Russia
 - 3.15 Spain
 - 3.16 Taiwan
 - 3.17 Turkey
 - 3.18 United Kingdom
 - 3.19 United States

⁴ See also



Comfort food

From Wikipedia, the free encyclopedia

For the Noah Ashenhurst novel, see [Comfort Food \(novel\)](#).

Comfort food is kinds of [food](#) that provides a [nostalgic](#) or sentimental value to someone,^[1] and may be characterized by its high [caloric nature](#), high [carbohydrate](#) level, or simple [preparation](#).^[2] The nostalgia may be specific to an individual, or it may apply to a specific [culture](#).^[3]



WIKIPEDIA
The Free Encyclopedia

Contents [hide]


- 1 Definition
- 2 Psychological studies
- 3 By region
 - 3.1 Afghanistan
 - 3.2 Australia and New Zealand
 - 3.3 Canada
 - 3.4 Egypt
 - 3.5 France
 - 3.6 Hong Kong
 - 3.7 India
 - 3.8 Indonesia
 - 3.9 Italy
 - 3.10 Japan
 - 3.11 Philippines
 - 3.12 Poland
 - 3.13 Puerto Rico
 - 3.14 Russia
 - 3.15 Spain
 - 3.16 Taiwan
 - 3.17 Turkey
 - 3.18 United Kingdom
 - 3.19 United States

Taiwan [edit]

Main article: [Taiwanese cuisine](#)

- Beef noodle soup^[79]
- Dan zai noodles — noodles and prawn with broth^[80]
- Minced pork rice^[80]
- Oyster omelette^[81]
- Red bean soup^[82]
- Small sausage in large sausage — grilled sausage in a rice casing^[81]
- Tangyuan — filled rice dumplings in sweet syrup^[82]

Brain-gut axis (2)



- Central serotonin production represents just 5% of total serotonin synthesis.
- The enterochromaffin cells in the gastrointestinal epithelium account for ~90% of all serotonin synthesis.
- It is becoming clear that the microbial influence on tryptophan metabolism and the serotonergic system

- “Complete” diet
 - Functional ingredients added



ROYAL CANIN VETERINARY DIET
Canine/Feline CALM dry food
(Unavailable in Taiwan)



Hill's Prescription Diet c/d
Multi-care Feline Stress



Hill's Prescription Diet b/d
Brain age care
(Unavailable in Taiwan)



Purina Pro Plan Veterinary Diets
NC NeuroCare
(Unavailable in Taiwan)

- Supplements
 - Complex mixtures
 - Combinations might be synergistic or antagonistic

成分個論

Classification of common dietary supplements ⁽³⁾

- Neuroprotective agent 神經保護
- Source of energy for the brain metabolism 參與神經代謝/提供能量
- Molecules of compounds that are structurally analogue to neurotransmitters 結構型似神經傳導物質
- Precursors of neurotransmitters 神經傳導物質的前驅物

L-Tryptophan(TRP) 左旋色胺酸

- Essential amino acid
- Foods that are rich in tryptophan: Chicken, banana, milk, soybean, chocolate...
- Precursors of neurotransmitters (serotonin and melatonin)
- 血清素系統(serotonergic system)低落容易發展成憂鬱或是焦慮傾向 (4, 5, 6)
 - 實驗動物：腦中serotonin低者對於社交壓力較敏感(4)
 - 狗：血中serotonin低者與表現攻擊性有關(5, 7)
 - 血清素本身無法通過血腦屏障



L-Tryptophan(TRP) 左旋色胺酸


- 可以穿過血腦障壁，但需與其他Large neural amino acid (LNAs)競爭⁽⁸⁾
- 需考量飲食中L-Tryptophan 和LNAs的比例，富含色胺酸的食物，通常也富含其它LNAs
- 無確實的建議補充劑量
- Single low-protein, high-carbohydrate diet led to a higher TRP/LNAs ratios and possibly, a higher TRP bioavailability in 6 hours.⁽⁷⁾
- 其他會影響色胺酸的因子⁽⁹⁾
 - Brain-gut axis⁽²⁾、Corticosteroids⁽⁸⁾、發炎、腸道菌叢、抗氧化物.etc.

L-Tryptophan(TRP) 左旋色胺酸

- 貓呢？

- Low-protein, high-carbohydrate diet for carnivores?
- The diet study (RC: Feline CALM dry food)^(10, 18)
 - Significantly increased the ratio of plasma tryptophan to large neutral amino acids
 - Decreased urinary cortisol concentrations
 - Multiple supplement : alpha-casozepine+ tryptophan

圖文不符之我是肉食獸...



L-Tryptophan(TRP) 左旋色胺酸

- Commonly found in commercial supplement mixture and nutraceutical diets.
- Generally calming/ anxiolytic effect.
 - Reduce stress-related behaviors in cats and working dogs⁽¹¹⁾
 - Reduce dominant and territorial aggression in dogs⁽¹²⁾
- Side effect induced by higher doses(70-220 mg/kg) were reported in human⁽¹³⁾.

L-Theanine 左旋茶胺酸

- Found in green tea
- Molecules of compounds that are structurally analogue to neurotransmitters
- Glutamate antagonist, 抑制興奮性的神經傳導物質
- 增加腦中特定部位的GABA, dopamine, serotonin
- 並無確實的建議劑量



L-Theanine 左旋茶胺酸

- Decrease social fear toward human⁽¹⁴⁾, and storm sensitivity in dogs⁽¹⁵⁾
- Decrease inappropriate toileting and avoidance behaviors in cats⁽¹⁶⁾
- Improvement takes 2-4 weeks to be seen, but twice a day for 2 months is recommended

Alpha-Casozepine

- Bovine milk protein alpha-S1 casein (酪蛋白) derivative
- Molecules of compounds that are structurally analogue to neurotransmitters (GABA_A agonist)
- Benzodiazepine-like effect
- Long-term treatment: 15mg/kg for 2-4 weeks
- Decrease fearful and anxious behaviors in cats ⁽¹⁰⁾
- Significantly decreased cortisol level in anxious dogs ⁽¹⁷⁾

Alpha-Lactalbumin

- Derived from the whey protein (乳清蛋白) of bovine milk
- Precursors of neurotransmitters (serotonin)
- Rich in amino acids including functional ingredients tryptophan
- Long-term treatment
- Improve cognitive tests and ability to cope with stress⁽¹¹⁾
- Serotonin depletions in brain caused by chronic anxiety

Melatonin 褪黑激素

- 內生性的激素
- 年紀增長，分泌漸不足
- Sleep-wake cycle disorders especially in elder animals
- Act as GABA_A agonist and may inhibit dopamine
- Anxiety, fear of thunder and fireworks
- 0.1mg/kg suggested (1.5-3 mg for small dogs/cats, 6-12 mg for large dogs)⁽¹⁹⁾
- 台灣未有保健品許可

S-adenosyl-L-methionine (SAMe)

- Neuroprotective agent/ Anti-oxidant
- Might increase serotonin levels, improve depressive state in human⁽¹⁹⁾
- Improve executive function in both aged dogs and cats⁽²⁰⁾
- Social interactions and disorientation in dogs with CDS⁽²¹⁾
- Should not be administration with tramadol⁽¹⁾

Medium Chain Triglycerides (MCT)

中鏈脂肪酸

- Source of energy for the brain metabolism
- 年紀增長，神經減少使用單醣作為能量來源
- MCT轉化為ketone bodies(酮類)作為替代能量來源
- Adjunctive treatment for cognitive dysfunction syndrome in dogs
- 須長期使用（一個月以上）
- 加入日常飲食，取代能量來源5%以上 (5.5%, 6.5%) (11)

Anti-oxidants 抗氧化物

- Vitamin: C, E
- Carotenoids 類胡蘿蔔素
- Ginkgo biloba 銀杏
- Reveratrol 白藜蘆醇
- Alpha-lipoic acid (硫辛酸) and L-carnitine(肉鹼)

國內外常見商品



Anxitane® S 安麗寧®

- L-Theanine
- 一天兩次，至少兩個月



Novifit® 蘇活沙美

- SAMe



Zylkene® (未正式上市)

- Alpha-Casozepine
- Social phobia in cats and anxiety in dogs
- Long-term and short-term effect



SOLLIQUIN® (未上市)

- L-Theanine/ Magnolia / Phellodendron/ Whey Protein (Alpha-Lactalbumin)
- Initial period for 30-45 days, or before event
- 一天一次，事件前可加強



TRANQUILE® 穩錠

- L-Tryptophan/ Taurine / Vitamin B3, 6, 12
- 長期使用：7天休息2天/ Before event



YuCALM® 優抗 犬/貓

- Lemon Balm / L-Theanine / Vitamin B complex/ Fish protein hydrolysate
- 一天一次



Nutrazyl® 樂寵壓易紓

- L-Tryptophan/ L-Theanine / Vitamin B complex/ GABA/ Passiflora



Composure®情緒調節保健 犬/貓嚼錠

- L-Theanine / Vitamin B1 complex/ Colostrum Calming Complex® Biopeptide Blend



紓壓胺基酸/情緒好放鬆保養粉

- Tryptophan/ L-Theanine / SAMe/ GABA/ 初乳蛋白



Senilife® 腦力活

- Phosphatidylserine/ pyridoxine/ ginko biloba extract/ resveratrol / Vit E
- Improve scores for disorientation, interaction, sleep-wake cycle and short-term memory in dogs⁽²²⁾

Golden Age 長青保®

- Acetyl-L-Carnitine/ Vitamin B complex/ Alpha lipoic acid/ Bioflavanol/N-Acetyl-Cysteine
- Anti-oxidant effects



Pet Remedy® 放輕鬆

- 繢草(Valeriana officinalis) 和其他草本萃取物
- GABAergic effects which work instantly



FELIWAY® (F3)

- Feline facial fraction F3
- Stress over resources and threats
- Transport and hospital stress

FELIFRIEND® (F4)


- Feline facial fraction F4
- Inter-cat aggression
- Aggression towards unfamiliar people





ADAPTIL® (未上市)

- Dog Appeasing Pheromones
- Anxiety and insecurity-related problems



ADAPTIL® Oral tablet (未上市)

- L-Tryptophan/ L-Theanine / GABA/ Vitamin B complex

注意事項

- Serotonin syndrome

- Neurological, autonomic, gastrointestinal symptoms: Seizures, tremors, depression, restlessness, mental confusion, tachycardia, hyperthermia, hypersalivation, GI conditions, coma, death...
- Avoid prescribing serotonergic medications and functional ingredients that also increase serotonin
- Serotonergic medications : SSRIs, SNAIs, TCAs, MOAIs...
- Functional ingredients : Tryptophan, SAMe, L-Theanine, Melatonin, Alpha-Lactalbumin...
- Risky combination : w Tramadol/ Fentanyl/ Amitraz/ Dextromethophan/ St. Johns wort/ 藤黃果/ 人蔘/ 紅景天...
- Prescription diet?



• Serotonin syndrome






Fig. 2. Pathways responsible for no, mild and severe serotonin syndrome

5-hydroxytryptophan (5-HTP)

- Toxicity in dogs, 90% (19/21) developed serotonin syndrome⁽²³⁾
- One capsule (500mg) would be sufficient to produce adverse effect in dogs.



感謝您的聆聽！
祝您情緒穩定！



有問題？請掃我！
Line ID: @827qbqot
E-mail: silent_vettw@gmail.com

References

1. Fernstrom JD. Large neutral amino acids: dietary effects on brain neurochemistry and function. *Amino Acids*. 2013 Sep;45(3):419-30.
2. Jenkins TA, Nguyen JC, Polglaze KE, Bertrand PP. Influence of Tryptophan and Serotonin on Mood and Cognition with a Possible Role of the Gut-Brain Axis. *Nutrients*. 2016 Jan 20;8(1):56D
3. <https://fabclinicians.org/wp-content/uploads/2020/11/OVERVIEW-OF-BEHAVIOUR-SUPPLEMENTS-FOR-DOGS-AND-CATS.pdf>
4. Sachs BD, Ni JR, Caron MG. Brain 5-HT deficiency increases stress vulnerability and impairs antidepressant responses following psychosocial stress. *Proc Natl Acad Sci U S A*. 2015 Feb 24;112(8):2557-62
5. Rosado B., Garcia-Belenguer S., Leon M., Chacon G., Villegas A., Palacio J. Blood concentrations of serotonin, cortisol and dehydroepiandrosterone in aggressive dogs. *Appl. Anim. Behav. Sci.* 2010;123:124–130
6. Ressler KJ, Nemeroff CB. Role of serotonergic and noradrenergic systems in the pathophysiology of depression and anxiety disorders. *Depress Anxiety*. 2000;12 Suppl 1:2-19..
7. Gazzano A, Ogi A, Torracca B, Mariti C, Casini L. Plasma Tryptophan/Large Neutral Amino Acids Ratio in Domestic Dogs Is Affected by a Single Meal with High Carbohydrates Level. *Animals (Basel)*. 2018;8(5):63. Published 2018 Apr 25
8. Dipace, V. Tryptophan effect on dog behavior: A review. *Dog Behavior*. 2015, 1, 23–31.
9. Strasser B, Gostner JM, Fuchs D. Mood, food, and cognition: role of tryptophan and serotonin. *Curr Opin Clin Nutr Metab Care*. 2016 Jan;19(1):55-61
10. Landsberg G, Milgram B, Mougeot I, Kelly S, de Rivera C. Therapeutic effects of an alpha-casozepine and L-tryptophan supplemented diet on fear and anxiety in the cat. *J Feline Med Surg*. 2017 Jun;19(6):594-602.
11. Orlando JM. Behavioral Nutraceuticals and Diets. *Vet Clin North Am Small Anim Pract*. 2018 May;48(3):473-495.
12. Jean S. DeNapoli, Nicholas H. Dodman, Louis Shuster, William M. Rand, and Kathy L. Gross. Effect of dietary protein content and tryptophan supplementation on dominance aggression, territorial aggression, and hyperactivity in dogs. *Journal of the American Veterinary Medical Association* 2000 217:4, 504-508
13. Crowell-Davis, Sharon & Murray, Thomas & Dantas, Leticia. (2019). *Veterinary Psychopharmacology*.
14. Araujo JA, De Rivera C, Ethier JE, Landsberg GM, et al. Anxitane tablets reduce fear of human beings in a laboratory model of anxiety-related behavior. *J Vet Behav*. 2010;5:268–75.
15. Pike AL, et al. An open-label prospective study of the use of L-theanine (Anxitane) in storm-sensitive client-owned dogs. *J Vet Behav: Clin Applications Res*. 2015;10(4):324–31.
16. Dramard, V., Kern, L., Hofmans, J. et al. Effect of L-theanine tablets in reducing stress-related emotional signs in cats: an open-label field study. *Ir Vet J* 71, 21 (2018)
17. Palestro, Clara & Minero, Michela & Cannas, Simona & Berteselli, Greta & Scaglia, Elisabetta & Barbieri, Sara & Cavallone, Elena & Puricelli, Maria & Servida, Francesco & Dall'Ara, Paola. (2010). Efficacy of a diet containing caseinate hydrolysate on signs of stress in dogs. *Journal of Veterinary Behavior-clinical Applications and Research - J VET BEHAV-CLIN APPL RES*. 5. 309-317.
18. Kazuki Miyaji, Maki Kato, Nobuyo Ohtani & Mitsuaki Ohta (2015) Experimental Verification of the Effects on Normal Domestic Cats by Feeding Prescription Diet for Decreasing Stress, *Journal of Applied Animal Welfare Science*, 18:4, 355-362
19. Overall, Karen. (2014). *Behavior Problems of the Dog and Cat*, 3rd ed., G. Landsberg, W. Hunthausen, L. Ackerman. Saunders, Elsevier (2013), 472
20. Araujo, Joseph & Faubert, Marjorie & Ba, & Brooks, Melissa & Landsberg, Gary & Lobprise, Heidi. NOVIFIT ® (NoviSAMe ®) Tablets Improve Executive Function in Aged Dogs and Cats: Implications for Treatment of Cognitive Dysfunction Syndrome. *Intern J Appl Res Vet Med* 2012; 10:90-8.
21. Mongillo, P., Araujo, J.A., de Rivera, C., Ethier, J.L., Landsberg, G.M., Bathurst, N., Denenberg, S., Milgram, N.W., 2010. The effects of Novifit on cognitive function in aged beagle dogs. *J. Vet. Behav.*
22. Osella M.C., Re G. ,Odore R. et al. Canine cognitive dysfunction syndrome: prevalence, clinical signs and treatment with a neuroprotective nutraceutical. *Appl Anim Behav Sci*. 2007; 105: 297-310
23. Gwaltney-Brant SM, Albretsen JC, Khan SA. 5-Hydroxytryptophan toxicosis in dogs: 21 cases (1989-1999). *J Am Vet Med Assoc*. 2000 Jun 15;216(12):1937-40